



Genorise[®] Recombinant Equine BMP4

Catalog Number: GR104020

Background

Bone morphogenetic protein 4 (BMP4) is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. This particular family member plays an important role in the onset of endochondral bone formation in humans. It has been shown to be involved in muscle development, bone mineralization, and uterine bud development. In human embryonic development, BMP4 is a critical signaling molecule required for the early differentiation of the embryo and establishing of a dorsal-ventral axis. BMP4 is secreted from the dorsal portion of the notochord, and it acts in concert with sonic hedgehog (released from the ventral portion of the notochord) to establish a dorsal-ventral axis for the differentiation of later structures. BMP4 stimulates differentiation of overlying ectodermal tissue. BMP4 has also been implicated in Fibrodysplasia Ossificans Progressiva in which it is overexpressed (1). Increase in expression of BMP4 has been associated with a variety of bone diseases, including the heritable disorder Fibrodysplasia Ossificans Progressiva.

References

1. Kan L, et al. (2004). *Am. J. Pathol.* 165 (4): 1107



Genorise® Recombinant Equine BMP4

Catalog Number: GR104020

Description

Size: 5 µg

Source: Chinese Hamster Ovary cell derived

Sequence: Ser293 –Arg409

Accession # NP_001157442.1

N-terminal Sequence Analysis: Ser293

Structure/Form: Disulfide-linked homodimer

Predicted Molecular Mass: 13 kDa (monomer)

Specifications

SDS-PAGE: 13 kDa, reducing conditions

Activity Measured in a cell proliferation assay using HUVEC human umbilical vein endothelial cells. Conn, G. et al. (1990) Proc Natl Acad Sci USA 87:1323. The ED50 for this effect is typically 15-38 ng/mL.

Endotoxin Level: <1.0 EU per 1 µg of the protein by the LAL method.

Purity: >96%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation: Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA with BSA as a carrier protein.

Preparation and Storage

Reconstitution: Reconstitute at 50 µg/mL in sterile PBS.

Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage: Use a manual defrost freezer and avoid repeated freeze thaw cycles.

- 6 months from date of receipt, -20 to -70°C as supplied.
- 1 month, -20 to -70°C under sterile conditions after reconstitution.

DECLARATION

THIS REAGENT IS FOR IN VITRO LABORATORY TESTING AND RESEARCH USE ONLY. DO NOT USE IT FOR CLINICAL DIAGNOSTICS. DO NOT USE OR INJECT IT IN HUMANS AND ANIMALS.