



Biotinylated Ovine IL-8 Polyclonal Antibody

Antigen-affinity purified donkey anti-ovine IL-8 antibody

Catalog Number: GR109106

Background

IL-8 was renamed CXCL8 by the Chemokine Nomenclature Subcommittee of the International Union of Immunological Societies (1), although its approved HUGO gene symbol remains IL8. CXCL8 was also referred to as neutrophil chemotactic factor (NCF), neutrophil activating protein (NAP), monocytederived neutrophil chemotactic factor (MDNCF), Tlymphocyte chemotactic factor (TCF), granulocyte chemotactic protein (GCP) and leukocyte adhesion inhibitor (LAI). Many cell types, including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, keratinocytes, hepatocytes, chondrocytes, and various tumor cell lines, can produce CXCL8 in response to a wide variety of proinflammatory stimuli such as exposure to IL1, TNF, LPS, and viruses. CXCL8 is a potent chemoattractant for neutrophils. In addition, CXCL8 also has a wide range of other proinflammatory effects. CXCL8 is also a potent angiogenic factor. CXCL8 causes degranulation of neutrophil specific granules and azurophilic granules. CXCL8 induces expression of the cell adhesion molecules CD11/CD18 and enhances the adherence of neutrophils to endothelial cells and subendothelial matrix proteins. Besides neutrophils, CXCL8 is also chemotactic for basophils, T cells and eosinophils. CXCL8 has been reported to be a comitogen for keratinocytes and was also shown to be an autocrine growth factor for melanoma cells. It is a key parameter in localized inflammation (2). It has been cited as a proinflammatory mediator in gingivitis and psoriasis (3).

References

1. Bacon K, et al (October 2002). *J. Interferon Cytokine Res.* 22 (10): 1067–8.
2. Vlahopoulos S, et al. (September 1999). *Blood* **94** (6): 1878–89.
3. Utgaard JO, et al (November 1998). *J. Exp. Med.* **188** (9): 1751–6.

Related products

1. GR238016 50 ml Reagent Reservoir, 100/case, 5 packs/case (pack of 20)
2. GR238004 Tissue Culture 96-well Microplate, individually packed, Case of 50
3. GR238019 1.5 ml Microcentrifuge tube with screw cap and free-standing, pack of 500
4. GR238007 125 ml leak-resistant HDPE bottle, colorless, pack of 24
5. GR238002 Microplate 12x8-Well Strip High Binding, Case of 50
6. GR238003 Microplate 12x8-Well Strip Medium Binding, Case of 50
7. GR238032 42592 Costar Stripwell Microplate 1 x 8 Flat Bottom, High Binding, Case of 100
8. GR238001 468667 Thermo Microplate 12x8-Well Strip Nunc Maxisorp F8, Case of 60
9. GR238031 96-well microplate sealer plastic, pack of 100



Biotinylated Ovine IL-8 Polyclonal Antibody

Antigen-affinity purified donkey anti-ovine IL-8 antibody

Catalog Number: GR109106

Description

Size: 50 µg

Species reactivity: Ovine, bovine

Specificity: Detects ovine IL-8 in direct ELISA and Western blot.

Source: Polyclonal Donkey IgG

Purification: Antigen-affinity purified

Immunogen: *E. coli* derived recombinant ovine IL-8, Ser29-Pro101, and Accession # NP_001009401.1

Endotoxin Level: < 0.10 EU per 1 µg of the antibody by the LAL method.

Conjugate: Biotin

Formulation: Lyophilized from 0.2 µm filtered PBS with 5% trehalose.

Application

Recommended concentrations:

Western blot: 0.2-0.5 µg/ml

Immunocytochemistry: 5-15 µg/ml

ELISA: 0.2-1.0 µg/ml

Neutralization: Measured by its ability to neutralize IL-8 induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CXCR2. The Neutralization Dose (ND₅₀) is typically 0.2-0.7 µg/mL in the presence of 20 ng/mL recombinant ovine IL-8.

Preparation, Stability & Storage

Reconstitution: Reconstitute at 0.2 mg/ml in sterile PBS with 0.05% azide.

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, at -20 °C to -70 °C, as supplied.
- 6 months from date of receipt, at -20 °C to -70 °C, after reconstitution.

DECLARATION

THIS REAGENT IS FOR IN VITRO LABORATORY TESTING AND RESEARCH USE ONLY. DO NOT USE IT FOR CLINICAL DIAGNOSTICS. DO NOT USE OR INJECT IT IN HUMANS AND ANIMALS.

**FOR LABORATORY RESEARCH USE ONLY
NOT FOR USE IN HUMANS AND ANIMALS**