



Genorise® Plasmid DNA Extraction Mini Kit 500

This kit uses mini-DNA binding column to isolate plasmid DNA from 1-5 ml of plasmid DNA-containing *E coli* and is for 500 applications. This kit can significantly increase quality and quantity of DNA and is more cost-effective than the similar products.

Materials provided in the kit:

1. 500 mini-DNA Binding columns
2. 500 Collection tubes
3. 300 ml Buffer GB
4. 150 ml Buffer GS
5. 150 ml Buffer GL
6. 200 ml Buffer GP
7. 140 ml Buffer GW1
8. 90 ml Buffer GW2
9. 60 ml Buffer GE

Materials required but not provided in the kit:

1. 100% Ethanol
2. 1.5 ml microcentrifuge tube

Reagent preparation

1. Buffer GW1: add 160 ml 100% ethanol
2. Buffer GW2: add 210 ml 100% ethanol

Protocol

1. Add 500 μ l Buffer GB to DNA binding column with collection tube, centrifuge 13,000 x g for 1 min and discard the flowthrough and keep the collection tube.
2. Centrifuge 1-5 ml bacterial culture at 13,000 x g for 1 min and discard supernatant. Pipet 250 μ l Buffer GS into pellet and suspend bacterial cells and transfer to a 1.5 ml microcentrifuge tube.
3. Add 250 μ l of Buffer GL and mix thoroughly by inverting the tube until solution becomes clear. Cell lysis should be done in 5 min.
4. Add 350 μ l Buffer GP and mix thoroughly by inverting the tube for 4-6 times. Centrifuge at 16,000 x g for 5 min.
5. Pipet the supernatant onto the column and with collection tube and centrifuge at full speed (20,000 x g) for 1 min. If samples remain in the column, repeat centrifugation to completely remove the remaining samples. Discard the flow-through and keep collection tube.
6. Add 500 μ l Buffer GW1 to the column in a new collection tube and add 500 μ l Buffer GW1. Centrifuge at 6000 x g for 1 min. Discard the flow-through and keep the collection tube.

7. Add 500 μ l Buffer GW2. Centrifuge at full speed (20,000 x g) for 3 min. Discard the flow-through and keep the collection tube.
8. Centrifuge at full speed for 1 min. Discard the flow-through and collection tube.
9. Place the mini column in a new 1.5 ml microcentrifuge tube and add 50-100 μ l Buffer GE and incubate for 2 min. Centrifuge at 6000 x g for 1 min to elute DNA.